


Pocket Reduction Surgery

INFORMATION
ABOUT POCKET
REDUCTION SURGERY





Pocket Reduction Surgery

0207 112 9036
info@rwperio.com
75 Harley St, Marylebone,
London W1G 8QL
www.rwperio.com

INFORMATION ABOUT POCKET REDUCTION SURGERY

What will it involve?

In some cases, non-surgical periodontal treatment does not produce enough of a reduction in the periodontal pocket. At this point, pocket reduction surgery may be suggested. If the pockets remain open and are not maintainable then the condition is likely to progress and may result in eventual tooth loss.

The procedure is carried out under local anaesthesia and you will be completely comfortable during the procedure. It will involve lifting the gums to gain direct access and full vision to the root surfaces. This will ensure thorough and predictable debridement. Furthermore, the gum and bone may be re-shaped and repositioned. Sometimes if the bone is of an appropriate shape a bone graft may be added (synthetic/human/animal-derived). If the latter is likely to be used, you will be informed beforehand. Once the procedure has been completed, the gums will be stitched back. You will need to come back 1-2 weeks later for the stitches to be removed and will be reviewed regularly thereafter.

Benefits

- Access to the roots of the teeth to allow for optimum debridement/ remove/cleaning.
- Ability to reshape the gums and bone.
- Ability to add bone grafts if appropriate.
- Reduction in the depth/closure of the periodontal pockets.
- Increased lifespan of the teeth.

Risks

- Pain can be managed using over-the-counter painkillers. These are usually required for 2-4 days.
- Swelling and bruising may occur depending on the length and extent of the procedure.
- A temporary increase in bleeding from the gums.
- Sensitivity but this is short-lasting and can be managed with toothpastes designed to reduce sensitivity.
- If the gums have been shaped, as they heal they may recede. This may mean your teeth look slightly longer and the spaces between them may be larger.
- Transient or permanent tooth looseness.

Timings

90 minutes for the surgery, 15-30 minutes for reviews.

Fees

Varies depending on complexity and number of teeth.

Examples:

RESHAPING THE GUMS

Before



After




ADDING A BONE GRAFT

Before



After



75 Harley St, Marylebone, London W1G 8QL
0207 112 9036 | info@rwperio.com
www.rwperio.com

